





27.02.2021, morning





A bit of fresh snow atop a hardened base

- 


Osterhorngruppe, Gamsfeldgruppe, Untersbergstock, Chiemgauer Alpen, Heutal, Reiteralpe, Steinernes Meer, Hochkönig, Hagengebirge, Göllstock, Loferer und Leoganger Steinberge, Tennengebirge, Gosaukamm


- 


Glocknergruppe Nord, Glocknergruppe Alpenhauptkamm, Goldberggruppe Nord, Goldberggruppe Alpenhauptkamm, Großvenedigergruppe Alpenhauptkamm, Großvenedigergruppe Nord, Oberpinzgauer Grasberge, Kitzbüheler Alpen, Glemmtal, Dientner Grasberge, Pongauer Grasberge, Niedere Tauern Alpenhauptkamm, Niedere Tauern Nord, Nockberge, Niedere Tauern Süd


- 

Ankogelgruppe, Muhr



Avalanche problems




Danger ratings



Expositions




27.02.2021, afternoon



Ein wenig Neuschnee auf harter Unterlage




Osterhorngruppe, Gamsfeld-Hinterglemm, Untersbergstock, Chiemgauer Alpen, Heutal, Reiteralpe, Steinernes Meer, Hochkönig, Hagengebirge, Göllstock, Loferer und Leoganger Steinberge, Tennengebirge, Gosaukamm



Glocknergruppe Nord, Glocknergruppe Alpenhauptkamm, Goldberggruppe Nord, Goldberggruppe Alpenhauptkamm, Großvenedigergruppe Alpenhauptkamm, Großvenedigergruppe Nord, Oberpinzgauer Grasberge, Kitzbüheler Alpen, Glemmtal, Dientner Grasberge, Pongauer Grasberge, Niedere Tauern Alpenhauptkamm, Niedere Tauern Nord, Nockberge, Niedere Tauern Süd




Ankogelgruppe, Muhr




2500 m


Avalanche problems



Danger ratings



Expositions



27.02.2021


Osterhorngruppe, Gamsfeldgruppe, Untersbergstock, Chiemgauer Alpen, Heutal, Reiteralpe, Steinernes Meer, Hochkönig, Hagengebirge, Göllstock, Loferer und Leoganger Steinberge, Tennengebirge, Gosaukamm



Stable powder snow on a hard base



Fresh snow, little wind



A small amount of fresh snow, noticeably lower temperatures

Avalanche danger is generally LOW. Due to solar radiation the fresh snow can naturally trigger small superficial loose-snow avalanches in extremely steep terrain. Small, in isolated cases medium-sized glide-snow avalanches continue to be possible in steep grass-covered terrain. Avalanche prone locations occur in isolated cases in extremely steep terrain on shady high altitude slopes.

Snowpack structure

Fresh snow (5-10 cm) will be deposited on a largely melt-freeze encrusted snowpack, on sunny slopes on bare ground up to 1600 m. The shallower-than-average snowpack has settled well, with good bonding to the layers beneath. On north-facing slopes the snowpack is still dry. In many zones the ground level depth hoar is the only weak layer.


Weather

On Saturday the peaks will often be veiled in heavy cloud, rainfall or snowfall in the early hours will not amount to much. The snowfall level will lie between 1000 and 1200 m. In the afternoon visibility will improve, clouds will disperse. A moderate northerly wind will be blowing. Lower temperatures than the last few days: at 1500 m; -3 degrees, at 2000 m, -5 degrees.


Outlook

On Sunday, sunny and a touch warmer, thus a slight daytime cycle of avalanche danger will return.

Avalanche problems



Danger ratings




Expositions



27.02.2021

Glocknergruppe Nord, Glocknergruppe Alpenhauptkamm, Goldberggruppe Nord, Goldberggruppe Alpenhauptkamm, Großvenedigergruppe Alpenhauptkamm, Großvenedigergruppe Nord, Oberpinzgauer Grasberge, Kitzbüheler Alpen, Glemmtal, Dientner Grasberge, Pongauer Grasberge, Niedere Tauern Alpenhauptkamm, Niedere Tauern Nord, Nockberge, Niedere Tauern Süd



very few danger zones in outlying terrain



on extremely steep grassy slopes, seldom and in extremely steep terrain

A few centimetres of fresh snow, noticeably lower temperatures

Avalanche danger is generally LOW. Due to solar radiation the fresh snow can naturally trigger small superficial loose-snow avalanches in extremely steep terrain. Small, in isolated cases medium-sized glide-snow avalanches continue to be possible in steep grass-covered terrain. Avalanche prone locations occur in isolated cases in extremely steep terrain on shady high altitude and high-alpine slopes.

Snowpack structure

A few centimetres of fresh snow will be deposited on a largely melt-freeze encrusted snowpack, on sunny slopes on bare ground up to 1600 m. The lower-than-average snowpack has settled well, with good bonding to the layers beneath. On north-facing slopes the snowpack is still dry. In many zones the ground level depth hoar is the only weak layer.

Weather

On Saturday the peaks will often be veiled in heavy cloud, rainfall or snowfall along and north of the Tauern in the early hours will not amount to much. The snowfall level will lie between 1000 and 1200 m. Peaks over 2000-2500 m might poke out above the fog. The mountains in Lungau will be favoured. In the afternoon visibility will improve, clouds will disperse. A moderate northerly wind will be blowing. Lower temperatures than the last few days: at 2000 m, -2 to -5 degrees, at 3000 m, -8 degrees.

Outlook

On Sunday, sunny weather, a touch warmer, thus a slight daytime cycle of naturally triggered avalanche danger will return.

Avalanche problems



New snow



Wind drifted snow



Persistent weak layer



Wet snow



Gliding snow



No problem

Danger ratings



1

low



2

moderate



3

considerable



4

high



5

very high

Expositions




27.02.2021, morning

Ankogelgruppe, Muhr



on extremely steep grassy slopes, possible at any time of day or night



A few centimetres of fresh snow, noticeably lower temperatures

Avalanche danger is generally LOW, rising during the daytime to MODERATE due to solar radiation (glide-snow problem). Small, in isolated cases medium-sized glide-snow avalanches continue to be possible in steep grass-covered terrain below 2500 m, particularly where glide cracks appear in the snowpack surface. Avalanche prone locations occur in isolated cases in extremely steep terrain on shady high altitude and high-alpine slopes.

Snowpack structure

Minor amounts of fresh snow will be deposited atop a melt-freeze encrusted snowpack surface (wind crusts, melt-freeze crusts), on sunny slopes the ground is often bare up to 1600 m. The old snowpack has settled well, generally is well bonded with the layers beneath. On north-facing slopes the snowpack is still dry. In many zones the ground-level soft layers of faceted crystals or depth hoar are the only weak layer. On steep sunny grass-covered slopes the entire snowpack can glide over the ground, in fact, is already gliding, though its movements have slowed. Despite lower temperatures, it has not stopped.


Weather

On Saturday morning, a few low-to-intermediate altitude residual clouds which will rapidly disperse. Sunshine will rapidly take over again. Temperatures will be measurably lower than recent days. At 2000 m: -2 to -5 degrees; at 3000 m, -8 degrees. At high altitudes, moderate northerly winds.


Outlook

On Sunday, sunny and slightly warmer, thus a slight daytime cycle of naturally triggered avalanche activity will return.

Avalanche problems



Danger ratings





Expositions




27.02.2021, afternoon

Ankogelgruppe, Muhr



  on extremely steep grassy slopes, possible at any time of day or night



A few centimetres of fresh snow, noticeably lower temperatures

Avalanche danger is generally LOW, rising during the daytime to MODERATE due to solar radiation (glide-snow problem). Small, in isolated cases medium-sized glide-snow avalanches continue to be possible in steep grass-covered terrain below 2500 m, particularly where glide cracks appear in the snowpack surface. Avalanche prone locations occur in isolated cases in extremely steep terrain on shady high altitude and high-alpine slopes.

Snowpack structure

Minor amounts of fresh snow will be deposited atop a melt-freeze encrusted snowpack surface (wind crusts, melt-freeze crusts), on sunny slopes the ground is often bare up to 1600 m. The old snowpack has settled well, generally is well bonded with the layers beneath. On north-facing slopes the snowpack is still dry. In many zones the ground-level soft layers of faceted crystals or depth hoar are the only weak layer. On steep sunny grass-covered slopes the entire snowpack can glide over the ground, in fact, is already gliding, though its movements have slowed. Despite lower temperatures, it has not stopped.

Weather

On Saturday morning, a few low-to-intermediate altitude residual clouds which will rapidly disperse. Sunshine will rapidly take over again. Temperatures will be measurably lower than recent days. At 2000 m: -2 to -5 degrees; at 3000 m, -8 degrees. At high altitudes, moderate northerly winds.

Outlook

On Sunday, sunny and slightly warmer, thus a slight daytime cycle of naturally triggered avalanche activity will return.

Translated by Jeffrey McCabe, www.creativtrans.com

Avalanche problems



Danger ratings



Expositions

