


Favorable conditions in the morning. Daytime increase of wet-snow avalanche danger.




Lechquellengebirge, Lechtaler Alpen, Verwall, Silvretta, Rätikon Ost, Rätikon West, Bregenzerwaldgebirge, Allgäuer Alpen



Avalanche problems




Danger ratings



Expositions





Vormittags günstige Bedingungen - mit der tageszeitlichen Erwärmung Anstieg der Gefahr von nassen Lawinen




Lechquellengebirge, Lechtaler Alpen, Verwall, Silvretta, Rätikon Ost, Rätikon West, Bregenzerwaldgebirge, Allgäuer Alpen



2400 m


Avalanche problems



Danger ratings




Expositions



Lechquellengebirge, Lechtaler Alpen, Verwall, Silvretta, Rätikon Ost, Rätikon West, Bregenzerwaldgebirge, Allgäuer Alpen



due to solar radiation and daytime warming



Slight increase in wet-snow avalanche danger below 2400m during daytime

Predominantly favorable conditions and low danger prevail in the morning. Due to solar radiation and daytime warming, avalanche danger increases slightly below 2400m. Naturally triggered or low-additional-loading triggered small-to-medium wet-snow avalanches occur especially on E/S/W facing slopes. Backcountry skiing tours and activities in outlying terrain should be launched and brought to an end early in the day. In regions where snowfall has been heavy on very steep grass-covered slopes, glide-snow avalanches continue to be possible. Caution below glide cracks. Weak layers in the old snow can often be triggered in high alpine regions, particularly by large additional loading small-to-medium avalanches can be triggered. Danger zones especially on very steep shady slopes.

Snowpack structure

Nighttime skies were partly cloudy, outgoing radiation was reduced compared to recent nights. Nevertheless, a crust formed which is generally capable of bearing loads. On sunny slopes this crust softens during the day due to solar radiation and daytime warming, forfeits its firmness, increasing the danger of wet-snow avalanches somewhat. At low and intermediate altitudes there is not much snow on the ground. In high alpine regions, particularly on very steep shady slopes, there are still isolated danger zones for dry-snow avalanches; small-to-medium slabs can trigger in the near-surface layers or at the Sahara dust layer.

Weather

Good mountain weather, some sunshine, some foehn. This afternoon some convective cloud build-up, these and other scattered clouds will not mar the pleasant weather much. At 2000 m: 0 to +5 degrees. Light southerly foehn wind, slightly intensifying later on.

Outlook

Due to the forecast higher temperatures and foehn influence, together with isolated rain showers, the snowpack will forfeit further its firmness. Avalanche danger could increase slightly from these factors.

Avalanche problems




Danger ratings




Expositions



Lechquellengebirge, Lechtaler Alpen, Verwall, Silvretta, Rätikon Ost, Rätikon West, Bregenzerwaldgebirge, Allgäuer Alpen



due to solar radiation and daytime warming



Slight increase in wet-snow avalanche danger below 2400m during daytime

Predominantly favorable conditions and low danger prevail in the morning. Due to solar radiation and daytime warming, avalanche danger increases slightly below 2400m. Naturally triggered or low-additional-loading triggered small-to-medium wet-snow avalanches occur especially on E/S/W facing slopes. Backcountry skiing tours and activities in outlying terrain should be launched and brought to an end early in the day. In regions where snowfall has been heavy on very steep grass-covered slopes, glide-snow avalanches continue to be possible. Caution below glide cracks. Weak layers in the old snow can often be triggered in high alpine regions, particularly by large additional loading small-to-medium avalanches can be triggered. Danger zones especially on very steep shady slopes.

Snowpack structure

Nighttime skies were partly cloudy, outgoing radiation was reduced compared to recent nights. Nevertheless, a crust formed which is generally capable of bearing loads. On sunny slopes this crust softens during the day due to solar radiation and daytime warming, forfeits its firmness, increasing the danger of wet-snow avalanches somewhat. At low and intermediate altitudes there is not much snow on the ground. In high alpine regions, particularly on very steep shady slopes, there are still isolated danger zones for dry-snow avalanches; small-to-medium slabs can trigger in the near-surface layers or at the Sahara dust layer.

Weather


Good mountain weather, some sunshine, some foehn. This afternoon some convective cloud build-up, these and other scattered clouds will not mar the pleasant weather much. At 2000 m: 0 to +5 degrees. Light southerly foehn wind, slightly intensifying later on.

Outlook


Due to the forecast higher temperatures and foehn influence, together with isolated rain showers, the snowpack will forfeit further its firmness. Avalanche danger could increase slightly from these factors.

Translated by Jeffrey McCabe, www.creativtrans.com

Avalanche problems



Danger ratings



Expositions

