


Avalanche report for Tuesday, 07.03.2023



Low avalanche danger in the Bavarian Alps



Bayerische Voralpen Mitte, Bayerische Voralpen Ost, Chiemgauer Alpen West, Chiemgauer Alpen Ost, Allgäuer Vorberge, Ammergauer Alpen, Werdenfeller Alpen, Bayerische Voralpen West, Allgäuer Hauptkamm, Berchtesgadener Alpen



Avalanche problems



Danger ratings






Expositions



Avalanche report for Tuesday, 07.03.2023

Bayerische Voralpen Mitte, Bayerische Voralpen Ost, Chiemgauer Alpen West, Chiemgauer Alpen Ost, Allgäuer Vorberge, Ammergauer Alpen, Werdenfeller Alpen, Bayerische Voralpen West, Allgäuer Hauptkamm, Berchtesgadener Alpen



Isolated small snowdrift accumulations are triggerable in high alpine regions - Attention to falls

Avalanche danger is low. Limited-space snowdrifts at heightened altitudes can be problematic. Small slab avalanches can be triggered by large additional loading near ridgelines on extremely steep shady slopes. Attention urged towards the risks of taking a fall, they outweigh those of being buried in snow masses.

In addition, on extremely steep sunny slopes at intermediate altitudes, small wet loose-snow avalanches can trigger naturally.

Snowpack structure

The snowpack has settled well overall, is largely stable. On sunny slopes the melt-freeze crust softens in the morning. At high altitudes the snowpack surface moistens only superficially. On shady slopes above 1700 m there is still powder. At high altitudes there are weak layers near small, older snowdrift accumulations.

Outlook

Over the next few days, danger of wet-snow avalanches at intermediate altitudes will increase. Snowdrift problem at high altitudes.

Translated by Jeffrey McCabe, www.creativtrans.com

Avalanche problems



Danger ratings



Expositions

